

## Comparison of aromatic composition of an endangered variety ('Albilla Dorada') with other recognized aromatic varieties

J. L. CHACÓN<sup>1)</sup>, E. GARCÍA<sup>1)</sup>, J. MARTÍNEZ<sup>1)</sup>, A. MENA<sup>1)</sup> and P. M. IZQUIERDO<sup>1),2)</sup>

<sup>1)</sup> Instituto de la Vid y el Vino de Castilla-La Mancha, Tomelloso, Spain

<sup>2)</sup> Parque Científico y Tecnológico de Albacete, Albacete, Spain

### Summary

'Albilla Dorada' is a local and endangered grape variety. It was correctly identified beforehand by analysing six microsatellite regions recommended by the GENRES 081 project plus six other additional regions, resulting in one new genotype not described for any variety according to the literature consulted. The object of this paper is to characterize the aromatic composition of this grape vine genotype not previously described. The minority volatile compounds in the grapes were analysed by gas chromatography-mass spectroscopy (GC-MS) and compared with those of eight known aromatic cultivars: 'Albillo Real', 'Macabeo', 'Malvasía Aromática', 'Gewürztraminer', 'Viognier', 'Malvasía Riojana', 'Riesling' and 'Moscatel de Grano Menudo'. Statistically significant differences between varieties were found as regards the different fractions making up the aromatic composition. 'Albilla Dorada' differed from the rest in that it had higher proportions of alcohols, benzene alcohols and norisoprenoids. In the case of the terpene fraction, it was the variety with the highest proportions of citronellol and terpene hydroxides. Results of this work point out 'Albilla Dorada' is a variety with a singular aromatic identity and highlight the necessity of recovering it before its total disappearance.

**Key words:** 'Albilla Dorada', genotype, grapes, volatile compounds.

### Introduction

The designation 'Albillo' or 'Albilla' (*Vitis vinifera* L.) is a generic term which includes a large group of minority white grape varieties grown in Spain under the same name. One of these is 'Albilla Dorada'. Recent studies have shown that this cultivar has a different genotype from all other previously-described 'Albillos' (FERNÁNDEZ-GONZÁLEZ *et al.* 2007).

'Albilla Dorada' is a local and endangered grape variety. It has been grown in the central Iberian Peninsula for many years and is in fact located exclusively in the municipal territory of Villamalea (La Mancha region). Its grapes are small and thin-skinned and acquire orange-pink tints upon ripening, which occurs in mid-August. The varietal

aroma is distinctive in every variety and is what contributes most to a wine's personality ('Moscatel', 'Chardonnay', 'Sauvignon Blanc', 'Riesling' ...). This is due essentially to the minority volatile compounds from the grape.

The object of this study is to determine the aromatic composition of 'Albilla Dorada' and see in what way it differs from that of other internationally-known aromatic varieties.

### Material and Methods

**Plant material:** This study was carried out in 2010. All grapes came from the same multivarietal vineyard, located in Tomelloso (La Mancha region-Spain). The following varieties were analysed: 'Albilla Dorada', 'Albillo Real', 'Macabeo', 'Malvasía Aromática', 'Gewürztraminer', 'Viognier', 'Malvasía Riojana', 'Riesling' and 'Moscatel de Grano Menudo' (or 'Muscat of Frontignan'). Varietal identity of the cultivars was established by analysing the twelve microsatellite regions: VVS2, VVMD5, VVMD7, VVMD27, VrZAG62, VrZAG79, VrZAG64, VrZAG67, VrZAG83, VVMD21, VVMD28 and VVMD36.

**Sample preparation and GC-MS analysis:** The GC-MS analytical method used was published by GARCÍA *et al.* (2003) and ROMERO *et al.* (2006). Samples were analysed in a TraceGCUltra gas chromatograph coupled to a DSQ mass spectrometer with an electron impact ionization source and a quadrupole analyser. For data processing, the minority volatile compounds were grouped into families by their chemical structure. To determine the concentrations of the minority volatile compounds, three samples of grapes of each variety were taken and processed separately.

### Results and Discussion

The genetic analysis showed 9 different profiles, eight of which belonged to known varieties. The genetic profile corresponding to 'Albilla Dorada' did not match the one proposed for any cultivar according to the data bases and literature consulted (FERNÁNDEZ-GONZÁLEZ *et al.* 2007, SANTANA *et al.* 2008).

Forty-six compounds were identified in the grapes: 8 C6 compounds, 4 linear alcohols, 18 terpene compounds (subdivided in terpenes, terpene oxides and terpene hy-






Figure: Projection of analysed samples onto the plane defined by the first two principal components of the percentages of the different volatile fractions.

jection of the samples analysed on to the plane defined by the two cited principal components. It was found that the sample distribution formed 3 groups based on the different percentages of the varietal aroma fractions.

A first group included the 'Moscatel de Grano Menudo' and 'Gewürztraminer' samples. These varieties are characterized first and foremost by a high percentage of terpene compounds and a smaller percentage of C<sub>6</sub> compounds than the rest, giving CP<sub>2</sub> values greater than zero. The second group, with negative CP<sub>2</sub> values and CP<sub>1</sub> values between -1 and 1, comprised the 'Albillo Real', 'Macabeo', 'Malvasía Aromática', 'Riesling', 'Malvasía Riojana' and 'Viognier' samples. The last group, consisting solely of

'Albilla Dorada', is substantially different from the others. All values of CP<sub>2</sub> are close to 0, and because of the high percentage of norisoprenoids, alcohols and benzene aldehydes and alcohols in this variety, it presents CP<sub>1</sub> values of between 2 and 3.

If we take only the terpene fraction, we find that as regards the monoterpenes in the grapes, the 'Albilla Dorada' variety (Table) contained a high percentage of citronellol (14.96 %) and a low percentage of Linalool (1.13 %). The highest proportion of complex forms was found in the form of hydroxides (75.49 %) which were considerably more abundant than in the other varieties.

Passport data on this cultivar can be found under "variety number vivc23429" in the Vitis International Variety Catalogue (<http://www.vivc.de>).

## References

- FERNÁNDEZ-GONZÁLEZ, M.; MENA, A.; IZQUIERDO, P.; MARTÍNEZ, J.; 2007: Genetic characterization of 142 grapevine (*Vitis vinifera* L.) cultivars from Castilla-La Mancha (Spain) using microsatellite markers. *Vitis* **46**, 126-131.
- GARCÍA, E.; CHACÓN, J. L.; MARTÍNEZ, J.; IZQUIERDO, P. M.; 2003: Changes in volatile compounds during ripening in grapes of Airen, Macabeo and Chardonnay white varieties grown in La Mancha Region (Spain). *Food Sci. Technol. Int.* **9**, 33-41.
- GÓMEZ, E.; MARTÍNEZ, A.; LAENCINA, J.; 1995: Changes in volatile compounds during maturation of some grape varieties. *J. Sci. Food Agric.* **67**, 229-233.
- MATTIVI, F.; GUZZON, R.; VRHOVSEK, U.; STEFANINI, M.; VELASCO, R.; 2006: Metabolite profiling of grape: Flavonols and anthocyanins. *J. Agric. Food Chem.* **54**, 7692-702.
- RODRÍGUEZ, R.; ROMERO, R.; CHACÓN, J. L.; MARTÍNEZ, J.; GARCÍA, E.; 2006: Phenolic compounds in skins and seeds of ten grape *Vitis vinifera* varieties grown in warm climate. *J. Food Comp. Anal.* **19**, 687-693.
- ROMERO, R.; CHACÓN, J. L.; GARCÍA, R.; MARTÍNEZ, J.; 2006: Pyrazine contents in four red grape varieties cultivated in a warm climate. *J. Int. Sci. Vigne Vin*, **40**, 203-207.
- SANTANA, J. C.; HIDALGO, E.; DE LUCAS, A. I.; RECIO, P.; ORTÍZ, J. M.; MARTÍN, J. P.; YUSTE, J.; ARRANZ, C.; RUBIO, J. A.; 2008: Identification and relationships of accessions grown in the grapevine (*Vitis vinifera* L.) Germplasm Bank of Castilla y León (Spain) and the varieties authorized in the VQPRD areas of the region by SSR-marker analysis. *Genet. Res. Crop Evol.* **55**, 573-583.

Received June 14, 2011

